




FELINE STRUVITE UROLITH DISSOLUTION

Diet plays a primary role in struvite management. Struvite stones dissolve in a few weeks and at a fraction of the cost of surgery. Kibble and canned formulations are effective. In five prospective studies with over 100 cats, dietary dissolution was 100 percent successful and no cats experienced urethral obstruction. Urethra stones should not be dissolved.

DISSOLUTION



*MN Urolith on iPhones and Minnesota Urolith Center on Androids

** We advise reviewing manufacturer's literature regarding selected therapeutic foods to determine indications and contraindications. For pets with multiple health concerns, we suggest that the selection of diet should take into consideration all health needs of the pet.

Support from [Hills Pet Nutrition](#), veterinarians, and pet owners make our work possible.





FELINE STRUVITE UROLITHS

Cats between 2 to 10 years old are at greatest risk for struvite urolith formation.¹ Although common in the urinary bladder, struvite is rare in the kidney. By reducing urinary phosphorus, magnesium, and pH, therapeutic foods are very effective at dissolving and preventing sterile struvite uroliths in cats.

Medical Considerations:

- Sterile struvite uroliths are common in cats. If urinary infection is considered, avoid empiric antimicrobics and culture urine first.
- Metabolic risk factors promoting alkalemia (renal tubular acidosis, hypoxemia, chronic diuretic use, administration of antacids, chronic vomiting, and hyperaldosteronism) and subsequent alkaluria are rarely diagnosed.

Nutritional Considerations:

- Diets restricted in phosphorus and magnesium that promote formation of acid urine (i.e. pH ≤ 6.4), minimize formation of sterile struvite uroliths.
- Extreme and prolonged reductions of some risk factors to minimize struvite urolith formation, including formation of acidic urine, may increase risk for calcium oxalate urolith formation. Therefore, we do not recommend long-term feeding of foods promoting urine pH < 6.2.¹
- High moisture foods (i.e. canned formulations) maybe more effective because increased water consumption is associated with decreased urine concentrations of calculogenic minerals and increased crystal evacuation. With persistent struvite crystalluria, feed canned foods and/or add increasing amounts of water to food until specific gravity is less than 1.030.

Pharmacological Considerations:

- Consider methionine or ammonium chloride to acidify the urine of patients consuming alternative diets that do not promote urine acidification.

Consider These Facts:

Some veterinarians prefer to remove struvite stones surgically instead of medical dissolution due to the perception that medical dissolution is less effective, more expensive, associated with prolonged discomfort, and will be associated with urethral obstruction as uroliths decrease in size.. These are misperceptions. Medical dissolution of sterile struvite uroliths was not associated with increased expense, urethral obstruction, or increased discomfort, and was more effective than surgery.^{2,3,4} Medical dissolution is an effective and compassionate choice for cats without a urethral obstruction.

Dissolution of sterile struvite uroliths with Prescription Diet[®] s/d[®] Feline pet food was 100% effective; in as quick as 6 days (range 6-28 days).⁴ Prescription Diet[®] c/d[®] Multicare Feline, a maintenance food specially formulated for long-term feeding, was also 100% successful in dissolving sterile struvite uroliths in as quick as 7 days (range 7 - 52 days).⁴ Consider feeding the maintenance food formulated to prevent struvite uroliths until cats are at least 9-10 years old and older if needed.¹

Struvite is the most common mineral in urethral plugs; following retropulsion or removal of urethra plugs feed therapeutic maintenance foods formulated to dissolve struvite.

Struvite crystalluria is common in cats with idiopathic cystitis. To eliminate risk for urethral obstruction, urolith formation, and bladder irritation from larger crystal aggregates; feed therapeutic maintenance foods formulated to dissolve and prevent struvite.

1. Lekcharoensuk C. Association between patient-related factors and risk of calcium oxalate and magnesium ammonium phosphate urolithiasis in cats: JAVMA 2000; 217: 520.

2. Osborne C. Medical dissolution of feline struvite urocystoliths: prospective clinical study of 30 cases. JAVMA 1990;196:1053.

3. Houston D. Evaluation of efficacy of a commercial diet in the dissolution of feline struvite bladder uroliths. Vet Ther 2004;5:187.

4. Lulich et al. Efficacy of two commercially available low magnesium, urine acidifying dry foods for the dissolution of struvite uroliths in cats. JAVMA 2013;243:1147.